arXiv:1411.5850v3 [math.CA] 7 Oct 2015

INEQUALITIES FOR DERIVATIVE OF POLYNOMIAL APPROXIMATION WITH ERDŐS TYPE WEIGHT

KENTARO ITOH, RYOZI SAKAI AND NORIAKI SUZUKI

1. Introduction

Let $\mathbb{R} = (-\infty, \infty)$. We consider an exponential weight

$$w(x) = \exp(-Q(x))$$

on $\mathbb{R}$, where Q is an even and nonnegative function on $\mathbb{R}$. Throughout this paper we always assume that w belongs to a relevant class $\mathcal{F}_\lambda(C^2+)$ (see Section 2). We consider a function $T = T_w$ defined by

$$T(x) := \frac{xQ'(x)}{Q(x)},\ x \neq 0 \tag{1.1}$$

is very important. We call w a Freud-type weight if T is bounded, and otherwise, w is called an Erdös-type weight.

For $x > 0$, the Mhaskar-Rakhmanov-Saff number (MRS number) $a_x = a_x(w)$ of $w = \exp(-Q)$ is defined by a positive root of the equation

$$x = \frac{2}{\pi}\int_0^1 \frac{a_x u Q'(a_x u)}{(1-u^2)^{1/2}}du. \tag{1.2}$$

When $w = \exp(-Q) \in \mathcal{F}(C^2+)$, Q' is positive and increasing on $(0, \infty)$, so that

$$\lim_{x\to\infty} a_x = \infty \quad \text{and} \quad \lim_{x\to\infty} \frac{a_x}{x} = 0 \tag{1.3}$$

hold. Note that those convergences are all monotonically.

For $1 \leq p \leq \infty$, we denote by $L^p(I)$ the usual L^p space on an interval I in $\mathbb{R}$. For $fw \in L^p(\mathbb{R})$ $(1 \leq p \leq \infty)$, the degree of weighted polynomial approximation $E_{p,n}(w, f)$ is defined by

$$E_{p,n}(w; f) := \inf_{P\in\mathcal{P}_n} \|w(f-P)\|_{L^p(\mathbb{R})},$$

where $\mathcal{P}_n$ is the class of all polynomials with degree n at most.

In [1], H. N. Mhaskar obtained the following result. Let $w = \exp(-Q)$ be a Freud-type weight, and let $1 \leq p \leq \infty$. Let f be absolutely continuous with $f'w \in L^p(\mathbb{R})$. There exists a constant $C > 0$ such that

2010 *Mathematics Subject Classification.* Primary 41A17, Secondary 41A10.

Key words and phrases. de la Vallée Poussin mean; Christoffel function; weighted polynomial approximation; Freud type weight; Erdős type weight.

for $n \in \mathbb{N}$ and $P \in \mathcal{P}_n$, if

$$\|w(f-P)\|_{L^p(\mathbb{R})} \le \eta$$

holds for $\eta > 0$, then

$$\|w(f'-P')\|_{L^p(\mathbb{R})} \le CE_{p,n-1}(w;f') + \frac{n}{a_n}\eta.$$

In particular, if we take $\eta = 2E_{p,n}(w,f)$, then

$$\|w(f'-P')\|_{L^p(\mathbb{R})} \le CE_{p,n-1}(w;f')$$

holds (see [1, Theorem 4.1.7]).

Now, we will obtain a counterpart of the above theorem for an Erdős-type weight. The following result is main theorem in this paper.

Theorem 1.1. Let $w = \exp(-Q)$ be a Erdős-type weight. Suppose that

$$T(a_n) \le c\left(\frac{n}{a_n}\right)^{2/3} \tag{1.4}$$

holds some constant $c \ge 0$. For any $C_1 \ge 1$, $1 \le p \le \infty$ and absolutely continuous function f with $f'w \in L^p(\mathbb{R})$, if

$$\|(f-P)w\|_{L^p(\mathbb{R})} \le C_1 E_{p,n}(w;f), \tag{1.5}$$

holds for some $n \in \mathbb{N}$ and $P \in \mathcal{P}_n$, then there exists constant $C_2 \ge 1$

$$\|w(f'-P')\|_{L^p(\mathbb{R})} \le C_2(1+C_1)T^{3/4}(a_n)E_{p,n-1}(w;f') \tag{1.6}$$

holds.

In particular, when we denote the n-degree best approximation polynomial of f with the weight w by $P_{f,n}$, that is,

$$E_{p,n}(w;f) = \|w(f-P)\|_{L^p}(\mathbb{R}), \tag{1.7}$$

then there exists a constant $C > 0$ such that

$$\|w(f'-P')\|_{L^p(\mathbb{R})} \le CT^{3/4}(a_n)E_{p,n-1}(w;f'). \tag{1.8}$$

Here if $p = \infty$, then we suppose that f' is continuous and $w(x)f'(x) \to 0$ as $|x| \to \infty$. When f' is continuous and $w(x)f'(x) \to 0$ as $|x| \to \infty$ we see $w(x)f(x) \to 0$ as $|x| \to \infty$. In fact, it follows from Lemma 2. 11.

Using (1.7) and (1.8), we have the following corollary.

Corollary 1.2. Let $w \in \mathcal{F}(C^2+)$ satisfy the condition (1.4). Let $1 \le p \le \infty$, for integer $r \ge 1$ and $n \ge 1$, there exists a constant $C_1 > 0$, and f^{r-1} be absolutely continuous in each compact interval with $f^{(r)}w \in L^p(\mathbb{R})$, where for $p = \infty$, we suppose that $f^{(r)}$ is continuous

and $w(x)f^{(r)}(x) \to 0$ as $|x| \to \infty$ (then we see $w(x)f^{(j)}(x) \to 0$ as $|x| \to \infty$, for $j = 0, 1, \cdots, r$). If for $P \in \mathcal{P}_n$, we have

$$\|w(f - P)\|_{L^p(\mathbb{R})} \le C_1 T^{1/4}(a_n) E_{p,n}(w; f),$$

then there exists a constant $C_2 > 0$ such that

$$\|w(f^{(j)} - P^{(j)})\|_{L^p(\mathbb{R})} \le C_2 T^{(2j+1)/4}(a_n) E_{p,n-j}(w; f^{(j)}).$$

2. class $\mathcal{F}(C^2+)$ and some fundamental notations

We say that an exponential weight $w = \exp(-Q)$ belongs to class $\mathcal{F}(C^2+)$, when $Q : \mathbb{R} \to [0, \infty)$ is a continuous and even function and satisfies the following conditions:
(a) $Q'(x)$ is continuous in $\mathbb{R}$ and $Q(0) = 0$.
(b) $Q''(x)$ exists and is positive in $\mathbb{R} \setminus \{0\}$.
(c) $\lim_{x\to\infty} Q(x) = \infty$.
(d) The function

$$T(x) := \frac{xQ'(x)}{Q(x)}, \ x \neq 0 \tag{1.1}$$

is quasi-increasing in $(0, \infty)$(i.e. there exists $C > 1$ such that $T(x) \le CT(y)$ whenever $0 < x < y$), and there exists $\Lambda \in \mathbb{R}$ such that

$$T(x) \ge \Lambda > 1, \ x \in \mathbb{R} \setminus \{0\}.$$

(e) There exists $C > 1$ such that

$$\frac{Q''(x)}{|Q'(x)|} \le C \frac{|Q'(x)|}{Q(x)}, \text{ a.e. } x \in \mathbb{R}.$$

Moreover, if there also exist a compact subinterval $J(\ni 0)$ of $\mathbb{R}$, and $C > 1$ such that

$$C\frac{Q''(x)}{|Q'(x)|} \ge \frac{|Q'(x)|}{Q(x)}, \text{ a.e. } x \in \mathbb{R} \setminus J.$$

Let $w \in \mathcal{F}(C^2+)$ and $\lambda > 0$. We write $w \in \mathcal{F}_\lambda(C^2+)$ if there exist $K > 1$ and $C > 1$ such that for all $|x| \ge K$,

$$\frac{|Q'(x)|}{Q(x)^\lambda} \le C$$

holds.

A typical example of Freud-type weight is $w(x) = \exp(-|x|^\alpha)$ with $\alpha > 1$. It belongs to $\mathcal{F}_\lambda(C^2+)$ for $\lambda = 1$. For $u \ge 0$, $\alpha > 0$ with $\alpha + u > 1$ and $l \in \mathbb{N}$, we set

$$Q(x) := |x|^u(\exp_l(|x|^\alpha) - \exp_l(0)),$$

where

$$\exp_l(x) := \exp(\exp(\exp(\cdots(\exp x)))) (l - \text{times}).$$

Then $w = \exp(-Q(x))$ is an Erdös-type weight, which belongs to $\mathcal{F}_\lambda(C^2+)$ for $\lambda > 1$.

Throughout this paper we always assume that w belongs to a class $\mathcal{F}(C^2+)$. Let $\{p_n\}$ be orthogonal polynomials for a weight w, that is, p_n is the polynomial of degree n such that

$$\int_{\mathbb{R}} p_n(x)p_m(x)w^2(x)dx = \delta_{mn}.$$

For a function f with $fw \in L^p(\mathbb{R})$, we set

$$s_n(f)(x) := \sum_{k=0}^{n-1} b_k(f)p_k(x) \text{ where } b_k(f) = \int_{\mathbb{R}} f(t)p_k(t)w^2(t)dt$$

for $n \in \mathbb{N}$ (the partial sum of Fourier series). Also it is given by

$$s_n(f)(x) = \int_{\mathbb{R}} K_n(x,t)f(t)w^2(t)dt,$$

where

$$K_n(x,t) = \sum_{k=0}^{n-1} p_k(x)p_k(t).$$

It is known that by the Cristoffel-Darboux formula

$$K_n(x,t) = \frac{\gamma_{n-1}}{\gamma_n}\frac{p_n(x)p_{n-1}(t) - p_n(t)p_{n-1}(x)}{x-t}$$

holds, where γ_n and γ_{n-1} are the leading coefficients of p_n and p_{n-1}, respectively. Then

$$a_n \sim \frac{\gamma_{n-1}}{\gamma_n}.$$

The de la Vallée Poussin mean $v_n(f)$ of f is defined by

$$v_n(f)(x) := \frac{1}{n}\sum_{j=n+1}^{2n} s_j(f)(x).$$

The Christoffel function $\lambda_n(x) = \lambda_n(w,x)$ is defined by

$$\lambda_n(x) := \frac{1}{K_n(x,x)} = \left(\sum_{k=0}^{n-1} p_k(x)^2\right)^{-1}.$$

Then

$$\lambda_n(x) = \inf_{P\in\mathcal{P}_{n-1}} \frac{1}{P(x)^2}\int_{\mathbb{R}} |P(t)w(t)|^2 dt.$$

holds on $\mathbb{R}$. We denote the zeros of the orthonormal polynomial $p_n(x)$ by $x_{n,n} < x_{n-1,n} < \cdots < x_{1,n}$. And then we define Christoffel numbers

$\lambda_{k,n},\ k = 1, 2, \ldots, n$ such as $\lambda_{k,n} := \lambda_n(x_{k,n})$. We also use the following function for $u > 0$:

$$\varphi_u(x) := \begin{cases} \dfrac{a_u}{u} \dfrac{1 - \frac{|x|}{a_{2u}}}{\sqrt{1 - \frac{|x|}{a_u} + \delta_u}}, & |x| \le a_u, \\ \\ \varphi_u(a_u), & a_u < |x|, \end{cases}$$

where $\delta_u := \{uT(a_u)\}^{-2/3}$.

The facts described below are very important.

Lemma 2.1. (1) [2, Lamma 3.5 (3.27)-(3.29)] For fixed $L > 0$ and $t > 0$,

$$a_{Lt} \sim a_t, \qquad T(a_{Lt}) \sim T(a_t),$$

and for $j = 0, 1$

$$Q^{(j)}(a_{Lt}) \sim Q^{(j)}(a_t).$$

(2) [2, Lamma 3.4 (3.17), (3.18)] For $t > 0$,

$$Q'(a_t) \sim \frac{t\sqrt{T(a_t)}}{a_t}, \qquad Q(a_t) \sim \frac{t}{\sqrt{T(a_t)}}.$$

(3) [2, Lamma 3.11] For fixed $L > 0$ and $t > 0$,

$$\left|1 - \frac{a_{Lt}}{a_t}\right| \sim \frac{1}{T(a_t)}.$$

Lemma 2.2. (a) [2, Corollary 13.4 (a) (12.20)] Uniformly for $n \ge 1$, $1 \le k \le n - 1$,

$$x_{k,n} - x_{k-1,n} \sim \varphi_n(x_{k,n}), \qquad \varphi_n(x_{k,n}) \sim \varphi_n(x_{k+1,n}).$$

(b) [3, Lemma 3.4 (d)] Let $\max|x_{k,n}|, |x_{k+1,n}| \le a_{n/2}$,

$$w(x_{k,n}) \sim w(x_{k+1,n}) \sim w(x), \quad (x_{k+1,n} \le x \le x_{k,n}).$$

So, for given $C > 0$ and $|x| \le a_{n/3}$, if $|x - x_{k,n}| \le C\varphi_n(x)$,

$$w(x) \sim w(x_{k,n}).$$

Lemma 2.3. [2, Theorem 9.3 (c)] Let $1 \le p \le \infty$ and $L > 0$. Then uniformly for $n \ge 1$ and $|x| \le a_n(1 + L\eta_n)$,

$$\lambda_n(x) \sim \varphi_n(x) w^p(x).$$

Moreover, uniformly for $x \in \mathbb{R}$,

$$\lambda_n(x) \ge C\varphi_n(x) w^p(x).$$

For any nonzero real valued functions $f(x)$ and $g(x)$, we write $f(x) \sim g(x)$ if there exist positive constants C_1 and C_2 independent of x, such that $C_1 g(x) \leq f(x) \leq C_2 g(x)$ holds for all x. Similarly, for any positive numbers $\{c_n\}_{n=1}^{\infty}$ and $\{d_n\}_{n=1}^{\infty}$, we write $c_n \sim d_n$.

Throughout this paper, $C_1, C_2, \ldots$ denote positive constants independent of n, x, t or polynomials $P_n(x)$. The same symbol does not necessarily denote the same constant in different occurrences. We consider the numbers p and q are $1 \leq p \leq \infty$ with $1/p + 1/q = 1$, if $p = \infty$ then $q = 1$ and $p = 1$ then $q = \infty$.

3. Lemmas

We prepare some important lemmas.

Lemma 3.1. [cf. 1, Appendeix A.1.1.] Let $fw \in L^p(\mathbb{R})$, then

$$E_{p,0}(w; f) = \sup_{\substack{\|gw\|_{L^q(\mathbb{R})} \leq 1 \\ \int_{\mathbb{R}} g(t) w^2(t) dt = 0}} \int_{\mathbb{R}} f(t) g(t) w^2(t) dt. \tag{3.1}$$

[Proof] For $fw \in L^p(\mathbb{R})$, $gw \in L^q(\mathbb{R})$ such that $\|gw\|_{L^q(\mathbb{R})} = 1$, $\int_{\mathbb{R}} g(t) w^2(t) dt = 0$ and any fixed $a \in \mathbb{R}$, we have

$$\begin{aligned} \left| \int_{\mathbb{R}} f(t) g(t) w^2(t) dt \right| &= \left| \int_{\mathbb{R}} (f(t) - a) g(t) w^2(t) dt \right| \\ &\leq \|(f - a)w\|_{L^p(\mathbb{R})} \|gw\|_{L^q(\mathbb{R})} \\ &= \|(f - a)w\|_{L^p(\mathbb{R})}. \end{aligned} \tag{3.2}$$

That is

$$E_{p,0}(w; f) \geq \sup_{\substack{\|gw\|_{L^q(\mathbb{R})} \leq 1 \\ \int_{\mathbb{R}} g(t) w^2(t) dt = 0}} \int_{\mathbb{R}} f(t) g(t) w^2(t) dt. \tag{3.3}$$

Next, we will proof the opposite direction. We first show the case of $1 < p < \infty$. For $fw \in L^p(\mathbb{R})$ and $c \in \mathbb{R}$, then $(f - c)w \in L^p(\mathbb{R})$. We put

$$\begin{aligned} I(c) :&= \int_{\mathbb{R}} \operatorname{sign} |f(t) - c|^{\frac{p}{q}} w^{\frac{p}{q}+1}(t) dt. \\ &= \int_{\mathbb{R}} \frac{(f(t) - c)}{|f(t) - c|} |f(t) - c|^{\frac{p}{q}} w^{\frac{p}{q}+1}(t) dt. \end{aligned}$$

Because $I(c)$ is continuous for every $c \in \mathbb{R}$ and $\lim_{c \to \infty} I(c) = -\infty$, $\lim_{c \to -\infty} I(c) = \infty$, so we can find a constant c_0 such that $I(c_0) = 0$. So we put

$$g(t) := \frac{(f(t) - c_0)|f(t) - c_0|^{\frac{p}{q}-1} w^{\frac{p}{q}-1}(t)}{\|(f - c_0)w\|_{L^p(\mathbb{R})}^{\frac{p}{q}}},$$

then

$$\|gw\|_{L^q(\mathbb{R})} = \left(\int_{\mathbb{R}} \frac{|f(t)-c_0|^p w^p(t)}{\|(f-c_0)w\|^p_{L^p(\mathbb{R})}} dt \right)^{\frac{1}{p}} = 1,$$

$$\begin{aligned} \int_{\mathbb{R}} g(t)w^2(t)dt &= \int_{\mathbb{R}} \operatorname{sign} \frac{|f(t)-c_0|^{\frac{p}{q}}}{\|(f-c_0)w\|^{\frac{p}{q}}} w^{\frac{p}{q}+1}(t)dt \\ &= \frac{I(c_0)}{\|(f-c_0)w\|^{\frac{p}{q}}} = 0 \end{aligned}$$

and

$$\begin{aligned} \int_{\mathbb{R}} f(t)g(t)w^2(t)dt &= \int_{\mathbb{R}} (f(t)-c_0)g(t)w^2(t)dt \\ &= \int_{\mathbb{R}} (f(t)-c_0) \frac{(f(t)-c_0)|f(t)-c_0|^{\frac{p}{q}-1} w^{\frac{p}{q}+1}}{\|(f-c_0)w\|^{\frac{p}{q}}_{L^p(\mathbb{R})}} dt \\ &= \frac{1}{\|(f-c_0)w\|^{\frac{p}{q}}_{L^p(\mathbb{R})}} \int_{\mathbb{R}} |f(t)-c_0|^{\frac{p}{q}+1} w^{\frac{p}{q}+1} dt \\ &= \frac{1}{\|(f-c_0)w\|^{\frac{p}{q}}_{L^p(\mathbb{R})}} \|(f-c_0)w\|^p_{L^p(\mathbb{R})} \\ &= \|(f-c_0)w\|_{L^p(\mathbb{R})}. \end{aligned}$$

That is,

$$E_{p,0}(w;f) \le \sup_{\substack{\|gw\|_{L^q(\mathbb{R})} \le 1 \\ \int_{\mathbb{R}} g(t)w^2(t)dt = 0}} \int_{\mathbb{R}} f(t)g(t)w^2(t)dt. \tag{3.4}$$

If $p = 1$, we put

$$\begin{aligned} I(c) :&= \int_{\mathbb{R}} \frac{(f(t)-c)}{|f(t)-c|} w(t)dt \\ &= \int_{\mathbb{R}} \operatorname{sign} w(t)dt. \end{aligned}$$

and let

$$c_1 := \sup_c \{I(c) > 0\}, \qquad c_2 := \inf_c \{I(c) < 0\}.$$

We note that $c_1 \le c_2$. If $c_1 < c_2$, we put $c_0 := (c_1+c_2)/2$, then $I(c_0) \le 0$ and $I(c_0) \ge 0$, that is $I(c_0) = 0$. On the other hand, if $c_1 = c_2 = c_0$, we put

$$\lim_{c \to c_0} I(c) = A, \qquad \lim_{c \to c_0} I(c) = B,$$

$$E_+ := \{x \in \mathbb{R}; f(x) - c_0 > 0\},$$
$$E_- := \{x \in \mathbb{R}; f(x) - c_0 < 0\},$$
$$E_0 := \{x \in \mathbb{R}; f(x) = c_0\}.$$

We may suppose $B < A$, then

$$\int_{E_0} w(t)dt > 0.$$

We note that

$$A = \int_{E_+ \cup E_0} w(t)dt - \int_{E_-} w(t)dt$$

and

$$B = \int_{E_+} w(t)dt - \int_{E_0 \cup E_-} w(t)dt.$$

So

$$A - B = 2\int_{E_0} w(t)dt.$$

Now we consider the case of $A \leq \int_{E_0} w(t)dt$ (tha case of $-B \leq \int_{E_0} w(t)dt$ is similarly). We put

$$g(t) := \frac{1}{w(t)} \times \begin{cases} 1 & t \in E_+ \\ 1 - A/(\int_{E_0} w(t)dt) & t \in E_0 \\ -1 & t \in E_- \end{cases}.$$

Then

$$\|gw\|_{L^\infty(\mathbb{R})} = 1$$

and

$$\begin{aligned} \int_{\mathbb{R}} g(t)w^2(t)dt &= \int_{E_+} w(t)dt + \int_{E_-} (-1)w(t)dt + \int_{E_0} w(t)g(t)dt \\ &= A - \int_{E_0} w(t)dt + \int_{E_0} w(t)g(t)dt \\ &= A - \int_{E_0} \frac{A}{\int_{E_0} w(t)dt} w(t)dt = 0, \end{aligned}$$

furthermore,

$$\begin{aligned}\int_{\mathbb{R}} f(t)g(t)w^2(t)dt &= \int_{\mathbb{R}} (f(t)-c_0)g(t)w^2(t)dt \\ &= \int_{E_+} (f(t)-c_0)w(t)dt - \int_{E_-} (f(t)-c_0)w(t)dt \\ &\quad + \int_{E_0} (f(t)-c_0)g(t)w^2(t)dt \\ &= \int_{E_+} |f(t)-c_0)|w(t)dt + \int_{E_-} |f(t)-c_0)|w(t)dt \\ &\quad + \int_{E_0} |f(t)-c_0)|w(t)dt \\ &= \|(f-c_0)w\|_{L^1(\mathbb{R})},\end{aligned}$$

That is,

$$E_{1,0}(w;f) \le \sup_{\substack{\|gw\|_{L^\infty(\mathbb{R})}\le 1 \\ \int_{\mathbb{R}} g(t)w^2(t)dt=0}} \int_{\mathbb{R}} f(t)g(t)w^2(t)dt. \tag{3.5}$$

Next, we show the case of $p=\infty$. For $fw \in L^\infty(\mathbb{R})$, we put

$$I(c) := \operatorname{ess\,inf}(f-c_0)w + \operatorname{ess\,sup}(f-c_0)w.$$

As the previous case, there exists a constant c_0 such that $I(c_0)=0$. For any small $\varepsilon>0$, let

$$\begin{aligned}F_+ &:= \left\{x\in\mathbb{R}; (f(x)-c_0)w(x) > \|(f-c_0)w\|_{L^\infty(\mathbb{R})} - \varepsilon\right\}, \\ F_- &:= \left\{x\in\mathbb{R}; (f(x)-c_0)w(x) < -\|(f-c_0)w\|_{L^\infty(\mathbb{R})} + \varepsilon\right\}.\end{aligned}$$

Then F_+ and F_- are disjoint. Let

$$A := \int_{\mathbb{R}} \chi_{F_+}(t)w^2(t)dt > 0, \quad B := \int_{\mathbb{R}} \chi_{F_-}(t)w^2(t)dt > 0$$

and

$$C := \frac{1}{2A}\int_{\mathbb{R}} \chi_{F_+}(t)w(t)dt + \frac{1}{2B}\int_{\mathbb{R}} \chi_{F_-}(t)w(t)dt.$$

Moreover,

$$g(t) := \frac{1}{C}\left(\frac{1}{2A}\chi_{F+}(t) - \frac{1}{2B}\chi_{F_-}(t)\right).$$

Then

$$\begin{aligned}\|gw\|_{L^1(\mathbb{R})} &= \int_{\mathbb{R}} \frac{1}{C}\left|\frac{1}{2A}\chi_{F+}(t) - \frac{1}{2B}\chi_{F_-}(t)\right| w(t)dt \\ &= \frac{1}{2C}\left(\frac{1}{A}\int_{\mathbb{R}} \chi_{F_+}(t)w(t)dt + \frac{1}{B}\int_{\mathbb{R}} \chi_{F_-}(t)w(t)dt\right) \\ &= 1\end{aligned}$$

and

$$\int_{\mathbb{R}} g(t)w^2(t)dt = C\left(\frac{1}{2A}\int_{\mathbb{R}} \chi_{F_+} w^2(t)dt - \frac{1}{2B}\int_{\mathbb{R}} \chi_{F_-} w^2(t)dt\right) = 0.$$

Moreover,

$$\begin{aligned}
\int_{\mathbb{R}} f(t)g(t)w^2(t)dt &= \int_{\mathbb{R}} (f(t)-c_0)g(t)w^2(t)dt \\
&= \int_{\mathbb{R}} C\left(\frac{(f(t)-c_0)}{2A}\chi_{F_+} - \frac{(f(t)-c_0)}{2B}\chi_{F_-}\right) w^2(t)dt \\
&= \int_{F_+} \frac{C(f(t)-c_0)}{2A} w^2(t)dt + \int_{F_-} \frac{-C(f(t)-c_0)}{2B} w^2(t)dt \\
&\geq \int_{F_+} \frac{C(\|(f-c_0)w\|_{L^\infty(\mathbb{R})} - \varepsilon)}{2A} w(t)dt \\
&\qquad + \int_{F_-} \frac{-C(\|(f-c_0)w\|_{L^\infty(\mathbb{R})} - \varepsilon)}{2B} w(t)dt \\
&= \|(f-c_0)w\|_{L^\infty(\mathbb{R})} - \varepsilon.
\end{aligned}$$

So we have

$$E_{\infty,0}(w;f) \leq \sup_{\substack{\|gw\|_{L^1(\mathbb{R})}\leq 1 \\ \int_{\mathbb{R}} g(t)w^2(t)dt=0}} \int_{\mathbb{R}} f(t)g(t)w^2(t)dt. \tag{3.6}$$

By (3.4), (3.5) and (3.6), the proof of the opposite direction is completed

Lemma 3.2 Let $P \in \mathcal{P}_n$. Then we have

$$\int_{\mathbb{R}} (f(t) - v_n(f)(t))P(t)w^2(t)dt = 0.$$

[Proof] Let $j \geq n+1$. For $P \in \mathcal{P}_n$, we have $P(t) = \sum_{k=0}^{n} a_k p_k(t)$. Therefore, we denote that $c_k := \int_{\mathbb{R}} f(x)p_k(x)w^2(x)dx$,

$$\begin{aligned}
&\int_{\mathbb{R}} (f(t) - s_j(f)(t))P(t)w^2(t)dt \\
&= \int_{\mathbb{R}} \left(f(t) - p(t)\sum_{k=0}^{j-1}\int_{\mathbb{R}} f(x)p_k(x)w^2(x)dx\right) P(t)w^2(t)dt \\
&= \int_{\mathbb{R}} \left(f(t) - \sum_{k=0}^{j-1} c_k p_k(t)\right) \sum_{l=0}^{n} a_l p_l(t)w^2(t)dt \\
&= \sum_{l=0}^{n} a_l c_l - \sum_{k=0}^{j-1} a_k c_k = 0.
\end{aligned}$$

Since $v_n(f) = \dfrac{1}{n}\displaystyle\sum_{j=n+1}^{2n} s_j(f)$, we have the result. $\square$

Lemma 3.3. For $x \in \mathbb{R}$ and $n \in \mathbb{N}$, we have

$$E_{1,2n-1}(w;\chi_x) \le C\frac{a_n}{n}w(x). \tag{3.7}$$

[Proof] Let $n \in \mathbb{N}$ be a fixed and integer $0 \le k \le n+1$ be found so that $x \in (x_{k+1,n}, x_{k,n}]$. Then there exist $P := P_k \in \mathcal{P}_{2n-2}$ and $R := R_k \in \mathcal{P}_{2n-2}$, such that

$$R_k(t) \le \chi_{(-\infty,x]}(t) \le P_k(t), \qquad (t \in \mathbb{R}) \tag{3.8}$$

and

$$\int_{\mathbb{R}} (P_k(t) - R_k(t))w^2(t)dt \le \lambda_{k+1,n} + \lambda_{k,n}. \tag{3.9}$$

(see [2, Corollary 1.2.6].). Let $|x| \le a_{n/3}$. First, we estimate $E_{1,n}(w^2;\chi_{(-\infty,x]})$. let k be an positive integer such tha $x \in [x_{k+1,n}, x_{k,n}] \subset [-a_n, a_n]$. Here, we note that $x_{1,n} < a_n$. There exist polynomials P and R satisfying (3.8) and (3.9), so that

$$\begin{aligned}
&E_{1,2n-1}(w^2;\chi_{(-\infty,x]}) \\
\le &\int_{\mathbb{R}} (P(t) - \chi_{(-\infty,x]}(t))w^2(t)dt + \int_{\mathbb{R}} (\chi_{(-\infty,x]}(t) - R(t))w^2(t)dt \\
\le &\lambda_{k+1,n} + \lambda_{k,n}.
\end{aligned}$$

By Lemma 2.3 (a), we have

$$\lambda_{k,n} \sim \varphi_n(x_{k,n})w^2(x_{k,n}),$$

and by Lemma 2.2 (b), if $|x - x_{k,n}| \le C\varphi_n(x)$, then $w(x) \sim w(x_{k,n})$. We also have $\varphi_n(x_{k,n}) \sim \varphi_n(x)$ by Lemma 2.2 (a). Therefore

$$\lambda_{k,n} \sim C\varphi_n(x)w^2(x).$$

Similarly, from $\varphi_n(x_{k+1}, n) \sim \varphi_n(x_{k,n}) \sim \varphi_n(x)$ and $|x - x_{k+1,n}| \le C\varphi_n(x)$, we have

$$\lambda_{k+1,n} \sim C\varphi_n(x)w^2(x).$$

Conseqently

$$E_{1,2n-1}(w^2;\chi_{(-\infty,x]}) \le C\varphi_n(x)w^2(x) \sim \frac{a_n}{n}\sqrt{1 - \frac{a_n}{n}}w^2(x) \le C\frac{a_n}{n}w^2(x).$$

Now, we estimate $E_{1,2n-1}(w;\chi_x)$. Since the MRS number for the weight w^2 equals a_{2n},

$$E_{1,2n-1}(w;\chi_x) = E_{1,2n-1}((w^{1/2})^2;\chi_{(-\infty,x]}) \le C\frac{a_{2n}}{n}w(x) \le C\frac{a_n}{n}w(x).$$

Next, we consider $|x| \geq a_{n/3}$. By Lemma 2.1 (2), we have

$$Q'(x) \geq Q'(a_{n/3}) \geq C\frac{(n/3)\sqrt{T(a_{n/3})}}{a_{n/3}} \geq C\frac{n\sqrt{T(a_n)}}{a_n}.$$

Suppose $x > 0$,

$$\begin{aligned} E_{1,2n-1}(w;\chi_{(-\infty,x]}) &\leq E_{1,0}(w;\chi_{(-\infty,x]}) \\ &\leq \int_{\mathbb{R}}(1-\chi_{(-\infty,x]}(t))w(t)dt = \int_x^\infty \exp(-Q(t))dt \\ &\leq \frac{-1}{Q'(x)}\int_x^\infty (-Q'(t))\exp(-Q(t))dt = \frac{w(x)}{Q'(x)} \\ &\leq C\frac{a_n}{n\sqrt{T(a_n)}}w(x) \leq C\frac{a_n}{n}w(x). \quad \square \end{aligned}$$

For $hw \in L^p(\mathbb{R})$, we set

$$I(h)(t) := \frac{1}{w^2(t)}\int_t^\infty h(u)w^2(u)du.$$

In following Lemma, weight w is Erdős type.

Lemma 3.4. (1) If $hw \in L^p(\mathbb{R})$ and

$$\int_{\mathbb{R}} h(t)w^2(t)dt = 0, \tag{3.10}$$

then

$$\|I'(h)w\|_{L^p(\mathbb{R})} \leq C\|hw\|_{L^p(\mathbb{R})} \tag{3.11}$$

Moreover, if g is absolutely continuous in each compact interval and $wg' \in L^q(\mathbb{R})$, then

$$\int_{\mathbb{R}} g(x)h(x)w^2(x)dx = \int_{\mathbb{R}} g'(t)I(h)(t)w^2(t)dt. \tag{3.12}$$

(2) If $hw \in L^\infty(\mathbb{R})$, and

$$\int_{\mathbb{R}} h(t)P(t)w^2(t)dt = 0, \quad P \in \mathcal{P}_n, \tag{3.13}$$

then

$$\|I(h)w\|_{L^\infty(\mathbb{R})} \leq C\frac{a_n}{n}\|hw\|_{L^\infty(\mathbb{R})}. \tag{3.14}$$

[Proof] We see that

$$\begin{aligned} I'(h)(t)w(t) &= \frac{2Q'(t)}{w(t)}\int_t^\infty w^2(u)h(u)du - h(t)w(t) \\ &= 2Q'(t)w(t)I(h)(t) - h(t)w(t). \end{aligned}$$

So, in order to prove (3.11), it is enough to show that

$$\|Q'I(h)w\|_{L^p(\mathbb{R})} \leq C\|hw\|_{L^p(\mathbb{R})}. \tag{3.15}$$

First we estimate for $p = \infty$. For $t > 0$, we have

$$\int_t^\infty w(x)dx \leq \int_t^\infty w(x)\left(1 + \frac{Q''(x)}{Q'^2(x)}\right)dx.$$

Since,

$$\frac{d}{dx}\left(\frac{w(x)}{Q'(x)}\right) = -w(x)\left(1 + \frac{Q''(x)}{Q'^2(x)}\right),$$

we have

$$\frac{w(t)}{Q'(t)} = \int_t^\infty w(x)\left(1 + \frac{Q''(x)}{Q'^2(x)}dx\right) \geq \int_t^\infty w(x)dx.$$

So, for all $t > 0$, we have

$$\frac{Q'(t)}{w(t)}\int_t^\infty w(x)dx \leq 1. \tag{3.16}$$

Therefore

$$\begin{aligned} |Q'(t)I(h)(t)w(t)| &= \left|\frac{Q'(t)}{w(t)}\int_t^\infty h(x)w^2(x)dx\right| \\ &\leq \|hw\|_{L^\infty(\mathbb{R})}\left|\frac{Q'(t)}{w(t)}\int_t^\infty w(x)dx\right| \\ &\leq \|hw\|_{L^\infty(\mathbb{R})}. \end{aligned} \tag{3.17}$$

If $t < 0$, (3.16) gives

$$\begin{aligned} \left|\frac{Q'(t)}{w(t)}\int_{-\infty}^t w(x)dx\right| &= \frac{Q'(|t|)}{w(|t|)}\int_{|t|}^\infty w(x)dx \\ &\leq 1. \end{aligned} \tag{3.18}$$

We may rewrite the condition (3.10) in the form

$$\int_t^\infty h(x)w^2(x)dx = -\int_{-\infty}^t h(x)w^2(x)dx,$$

so by (3.18), for $t < 0$,

$$\begin{aligned} \left|\frac{Q'(t)}{w(t)}\int_t^\infty h(x)w^2(x)dx\right| &= \left|\frac{Q'(t)}{w(t)}\int_{-\infty}^t h(x)w^2(x)dx\right| \\ &\leq \|hw\|_{L^\infty(\mathbb{R})}\left|\frac{Q'(t)}{w(t)}\int_{|t|}^\infty w(x)dx\right| \\ &\leq \|hw\|_{L^\infty(\mathbb{R})}. \end{aligned}$$

Therefore, hor all h satisfying (3.10), we have shown that

$$\|Q'I(h)w\|_{L^\infty(\mathbb{R})} \leq \|hw\|_{L^\infty(\mathbb{R})}. \tag{3.19}$$

Next, we estimate for $p = 1$. Let $H(x) := |h(x)| + |h(-x)|$, we have

$$\begin{aligned}
&\|Q'I(h)w\|_{L^1(\mathbb{R})} = \int_{\mathbb{R}} \left| \frac{Q'(t)}{w(t)} \int_t^\infty h(x)w^2(x)dx \right| dt \\
&= \int_0^\infty \left| \frac{Q'(t)}{w(t)} \int_t^\infty h(x)w^2(x)dx \right| dt + \int_{-\infty}^0 \left| \frac{Q'(t)}{w(t)} \int_t^\infty h(x)w^2(x)dx \right| dt \\
&= \int_0^\infty \left| \frac{Q'(t)}{w(t)} \int_t^\infty h(x)w^2(x)dx \right| dt + \int_{-\infty}^0 \left| \frac{Q'(t)}{w(t)} \int_{-\infty}^t h(x)w^2(x)dx \right| dt \\
&= \int_0^\infty \left| \frac{Q'(t)}{w(t)} \int_t^\infty h(x)w^2(x)dx \right| dt + \int_0^\infty \left| \frac{Q'(s)}{w(s)} \int_{-\infty}^{-s} h(x)w^2(x)dx \right| ds \\
&\le \int_0^\infty \frac{Q'(t)}{w(t)} \left(\int_t^\infty H(x)w^2(x)dx \right) dt = \int_0^\infty H(x)w^2(x) \left(\int_0^x \frac{Q'(t)}{w(t)} dt \right) dx \\
&= \int_0^\infty H(x)w^2(x) \left(\frac{1}{w(x)} - 1 \right) dx \le \int_0^\infty H(x)w^2(x) \frac{1}{w(x)} dx \\
&= \int_{\mathbb{R}} |h(x)|w(x)dx = \|hw\|_{L^1(\mathbb{R})}.
\end{aligned} \tag{3.20}$$

Now, let $fw \in L^p(\mathbb{R})(1 \le p \le \infty)$. We define a linear operator

$$F(f) := \frac{\displaystyle\int_{\mathbb{R}} f(x)w^2(x)dx}{\displaystyle\int_{\mathbb{R}} w^2(x)dx}.$$

Then we have

$$|F(f)| \le C\|fw\|_{L^p(\mathbb{R})} \tag{3.21}$$

for $p = 1, \infty$. And so

$$\|(f - F(f))w\|_{L^p(\mathbb{R})} \le C\|fw\|_{L^p(\mathbb{R})} \tag{3.22}$$

for $p = 1, \infty$, that is $(f - F(f))w \in L^p(\mathbb{R})$ for $p = 1, \infty$. Next, we have

$$\begin{aligned}
\int_{\mathbb{R}} (f(t) - F(f))w^2(t)dt &= \int_{\mathbb{R}} f(t)w^2(t)dt - F(f) \int_{\mathbb{R}} w^2(t)dt \\
&= \int_{\mathbb{R}} f(t)w^2(t)dt - \int_{\mathbb{R}} f(x)w^2(x)dx = 0,
\end{aligned}$$

that is $h := f - F(f)$ satisfies (3.10). Let $p = 1, \infty$ and $fw \in L^p(\mathbb{R})$, we consider the operator

$$U(f) := Q'I(f - F(f)).$$

Then, from (3.17), (3.20) and (3.22), we conclude that

$$\|U(f)w\|_{L^p(\mathbb{R})} \le C\|(f - F(f))w\|_{L^p(\mathbb{R})} \le C\|fw\|_{L^p(\mathbb{R})} \tag{3.23}$$

By Riesz-Thorin interpolation theorem, the estimate (2.28) holds for all $1 \le p \le \infty$. Therefore

$$\|U(f)w\|_{L^p(\mathbb{R})} = \|Q'I(f - F(f))w\|_{L^p(\mathbb{R})} \le C\|fw\|_{L^p(\mathbb{R})}. \tag{3.24}$$

Now we see that if h satisfies (3.10), then we have $F(h) = 0$. So applying (3.24) to $f = h$, we conclude

$$\|Q'I(h)w\|_{L^p(\mathbb{R})} \le C\|hw\|_{L^p(\mathbb{R})}$$

for all $1 \le p \le \infty$. This completes the proof of (3.15), that is (3.11).

Next, for $1 < p < \infty$, in the same way as (3.18), for $t < 0$, we have

$$|qQ'(t)| \int_{-\infty}^{t} w^q(x)dx \le w^q(t)$$

and since w is Erdős type, we have

$$\frac{1}{|Q'(t)|} \le C\frac{1}{|t|^{2q/p}}$$

holds (see [4, (2.4)]). Now let $g'w \in L^q(\mathbb{R})$, then $gw \in L^q(\mathbb{R})$ (cf. [4, Theorem 6]), hence

$$\begin{aligned}
&\int_{-\infty}^{-1} |g'(t)| \left(\int_{-\infty}^{t} |h(x)|w^2(x)dx \right) dt \\
&\le \int_{-\infty}^{-1} |g'(t)| \left(\int_{-\infty}^{t} (|h(x)|w(x))^p dx \right)^{1/p} \left(\int_{-\infty}^{t} w^q(x)dx \right)^{1/q} dt \\
&\le \|hw\|_{L^p(\mathbb{R})} \int_{-\infty}^{-1} |g'(t)| \left(\frac{w^q(t)}{Q'(t)} \right)^{1/q} dt \\
&= \|hw\|_{L^p(\mathbb{R})} \int_{-\infty}^{-1} |g'(t)|w(t) \frac{1}{(Q'(t))^{1/q}} dt \\
&\le \|hw\|_{L^p(\mathbb{R})} \|g'w\|_{L^q(\mathbb{R})} \left(\int_{-\infty}^{-1} \frac{1}{t^2} dt \right)^{1/p} < \infty.
\end{aligned}$$

If $p = \infty$, then

$$\begin{aligned}
&\int_{-\infty}^{-1} |g'(t)| \left(\int_{-\infty}^{t} |h(x)|w^2(x)dx \right) dt \\
&\le \int_{-\infty}^{-1} \left(|g'(t)| \|hw\|_{L^\infty(\mathbb{R})} \int_{-\infty}^{t} w(x)dx \right) dt \\
&\le \|hw\|_{L^\infty(\mathbb{R})} \int_{-\infty}^{-1} \frac{w(t)}{|Q'(t)|} |g'(t)| dt \\
&\le C\|hw\|_{L^\infty(\mathbb{R})} \int_{-\infty}^{-1} |g'(t)|w(t)dt < \infty.
\end{aligned}$$

Finally, we consider $p = 1$, then

$$\begin{aligned}
&\int_{-\infty}^{-1} |g'(t)| \left(\int_{-\infty}^{t} |h(x)| w^2(x) dx \right) dt \\
&= \int_{-\infty}^{-1} |h(x)| w^2(x) \left(\int_{x}^{-1} |g'(t)| dt \right) dx \\
&= \int_{-\infty}^{-1} |h(x)| w(x) \left(\int_{x}^{-1} |g'(t)| w(t) \frac{w(x)}{w(t)} dt \right) dx.
\end{aligned}$$

We note that

$$\int_x^{-1} \frac{w(x)}{w(t)} dt = w(x) \int_x^{-1} \frac{Q'(t)}{w(t)} \frac{1}{Q'(t)} dt \leq \frac{w(x)}{C} \int_x^{-1} \frac{Q'(x)}{w(t)} dt < \frac{1}{C}.$$

holds. So

$$\int_{-\infty}^{-1} |g'(t)| \left(\int_{-\infty}^{t} |h(x)| w^2(x) dx \right) dt < \infty.$$

And

$$\int_{-1}^{0} |g'(t)| \left(\int_{-\infty}^{t} |h(x)| w^2(x) dx \right) dt < \infty.$$

By (3.10), we may assume thet $g(0) = 0$ in (3.12). Then

$$\begin{aligned}
&\int_{-\infty}^{\infty} g(x) h(x) w^2(x) dx \\
&= -\int_{-\infty}^{0} h(x) w^2(x) \left(\int_{x}^{0} g'(t) dt \right) dx \\
&= -\int_{-\infty}^{0} \left(\int_{-\infty}^{0} g'(t) \chi_{[x,0]}(t) dt \right) h(x) w^2(x) dx \\
&= -\int_{-\infty}^{0} \left(\int_{-\infty}^{0} h(x) \chi_{(-\infty,t]}(x) w^2(x) dx \right) g'(t) dt \\
&= -\int_{-\infty}^{0} g'(t) \left(\int_{-\infty}^{t} h(x) w^2(x) dx \right) dt \\
&= \int_{-\infty}^{0} g'(t) \left(\int_{t}^{\infty} h(x) w^2(x) dx \right) dt \\
&= \int_{-\infty}^{0} g'(t) I(h)(t) w^2(t) dt.
\end{aligned}$$

Similarly,

$$\int_0^{\infty} g(x) h(x) w^2(x) dx = \int_0^{\infty} g'(t) I(h,t) w^2(t) dt.$$

Hence, (3.12) is proved.

Moreover, let h satisfies (3.13), we get for $t \in \mathbb{R}$ and $P_{\chi,t} \in \mathcal{P}_{2n-1}$ is the best approximation for $\chi_{[-\infty,t]}$ in $L^1(\mathbb{R})$. Then

$$\begin{aligned}
|I(h)(t)| &= \left| \frac{1}{w^2(t)} \int_t^\infty h(x)w^2(x)dx \right| \\
&= \left| \frac{1}{w^2(t)} \int_{-\infty}^t h(x)w^2(x)dx \right| \\
&= \left| \frac{1}{w^2(t)} \int_{\mathbb{R}} \chi_{(-\infty,t]}(x)h(x)w^2(x)dx \right| \\
&= \left| \frac{1}{w^2(t)} \int_{\mathbb{R}} (\chi_{(-\infty,t]}(x) - P_{\chi,t}(x))h(x)w^2(x)dx \right| \\
&\leq \frac{\|hw\|_{L^\infty(\mathbb{R})}}{w^2(t)} \int_{\mathbb{R}} |\chi_{(-\infty,t]}(x) - P_{\chi,t}(x)|w(x)dx \\
&\leq \frac{\|hw\|_{L^\infty(\mathbb{R})}}{w^2(t)} E_{1,2n-1}(w; \chi_{(-\infty,t]}) \\
&\leq C\frac{1}{w(t)}\frac{a_n}{n}\|hw\|_{L^\infty(\mathbb{R})},
\end{aligned}$$

By Lemma 3.3. Therefore

$$\|I(h,t)w\|_{L^\infty(\mathbb{R})} \leq C\frac{a_n}{n}\|hw\|_{L^\infty(\mathbb{R})}. \quad \square$$

Lemma 3.5. [4, Theorem 1]. Let g be absolutely continuous in each compact interval and $g'w \in L^p(\mathbb{R})$. Then we have

$$E_{p,n}(w; g) \leq C\frac{a_n}{n}\|g'w\|_{L^p(\mathbb{R})},$$

and equivalently,

$$E_{p,n}(w; g) \leq C\frac{a_n}{n}E_{p,n-1}(w; g').$$

Lemma 3.6. [5. Theorem 1.1]. Let $1 \leq p \leq \infty$ and $T(a_n) \leq c(n/a_n)^{2/3}$ for some $c > 0$. If $T^{1/4}fw \in L^p(\mathbb{R})$, then

$$\|(f - v_n(f))w\|_{L^p(\mathbb{R})} \leq CT^{1/4}(a_n)E_{p,n}(w; f).$$

Lemma 3.7. Let g be absolutely continuous and $g' \in L^p(\mathbb{R})$, $T(a_n) \leq c(n/a_n)^{2/3}$ for some $c > 0$.. Then for $n \in \mathbb{N}$ there exists a polynomial $V_n(g) \in \mathcal{P}_{2n}$ such that $V_n' = v_n(g')$ and

$$\|(g - V_n)w\|_{L^p(\mathbb{R})} \leq C\frac{a_n}{n}T^{1/4}(a_n)E_{p,n}(w; g'). \tag{3.25}$$

[Proof] Without loss of generality, we may assume the $g(0) = 0$. Let

$$G(x) := \int_0^x (g'(t) - v_n(g')(t))dt,$$

and take a constant a such that

$$\|(G-a)w\|_{L^p(\mathbb{R})} \le 2E_{p,0}(w;G).$$

Then

$$\begin{aligned} V_n(x) : &= a + \int_0^x v_n(g')(t)dt \\ &= a + g(x) - G(x) \in \mathcal{P}_{2n}, \end{aligned} \tag{3.26}$$

and

$$\|(g-V_n)w\|_{L^p(\mathbb{R})} = \|(G-a)w\|_{L^p(\mathbb{R})} \le 2E_{p,0}(w;G). \tag{3.27}$$

By the duality principle and Lemma 3.1, we find h such thae $hw \in L^q(\mathbb{R})$ and satisfies (3.1), $\|hw\|_{L^q(\mathbb{R})} = 1$ and by Lamma 3.2, in the same way as changing the order of integration in proof of (3.12),

$$\begin{aligned} &\|(g - V_n(g))w\|_{L^p(\mathbb{R})} \le 2E_{p,0}(w;G) \\ &\le 4\left|\int_{\mathbb{R}} G(x)h(x)w^2(x)dx\right| \\ &= 4\int_{\mathbb{R}} \left(\int_0^x g'(t) - v_n(g')(t)dt\right) h(x)w^2(x)dx \\ &= 4\int_{\mathbb{R}} (g'(t) - v_n(g')(t)) \left(\int_t^\infty h(x)w^2(x)dx\right) dt \\ &= 4\int_{\mathbb{R}} (g'(t) - v_n(g')(t))I(h)(t)w^2(t)dt \\ &= 4\int_{\mathbb{R}} (g'(t) - v_n(g')(t))(I(h)(t) - P_{I(h)})w^2(t)dt \end{aligned}$$

holds(cf. [4. Theorem 6]). Where $P_{I(h)} \in \mathcal{P}_n$ is the best approximation polynomial for $I(h)$ with $L^q(\mathbb{R})$. From Lemma 3.5 with q and Lemma 3.4 (3.11) with q, we have

$$E_{q,n}(w;I(h)) \le C\frac{a_n}{n}\|I'(h)w\|_{L^q(\mathbb{R})} \le C\frac{a_n}{n}\|hw\|_{L^q(\mathbb{R})} = C\frac{a_n}{n}. \tag{3.28}$$

Since by Lemma 3.6,

$$\|(g' - v_n(g')w\|_{L^p(\mathbb{R})} \le CT^{1/4}(a_n)E_{p,n}(w;g'). \tag{3.29}$$

From (3.28) and (3.29), we have the result. $\square$.

4. Proof of Theorem 1.1

Theorem 4.1. Let $w \in \mathcal{F}(C^2+)$ is Erdős type and suppose that $T(a_n) \le c(n/a_n)^{2/3}$ holds some constant $c > 0$. Then there exists a constant $C \ge 1$ such that for any absolutely continuous function f with $f'w \in L^q(\mathbb{R})$, if

$$\|(f-P)w\|_{L^p(\mathbb{R})} \le \eta$$

holds for some $n \in \mathbb{N}$, $P \in \mathcal{P}_n$ and $\eta > 0$, then

$$\|w(f'-P')\|_{L^p(\mathbb{R})} \le C\left(T^{3/4}(a_n)E_{p,n}(w;f') + \frac{n}{a_n}T^{1/2}(a_n)\eta\right). \tag{4.1}$$

[Proof] By [2, Theorem 1.15 and Corollary 1.16], we see

$$\|P'w\|_{L^p(\mathbb{R})} \le C\frac{nT^{1/2}(a_n)}{a_n}\|Pw\|_{L^p(\mathbb{R})}$$

and by Proposition 3.3, we have

$$\|(f-V_n)w\|_{L^p(\mathbb{R})} \le C\frac{a_n}{n}T^{1/4}(a_n)E_{p,n}(w;f')$$

where $V_n' = v_n(f')$. Also by (3.29), we have

$$\|(f'-v_n(f'))w\|_{L^p(\mathbb{R})} \le CT^{1/4}(a_n)E_{p,n}(w;f'). \tag{4.2}$$

Hence

$$\begin{aligned}
&\|(f'-P')w\|_{L^p(\mathbb{R})} \le \|(f'-v_n(f'))w\|_{L^p(\mathbb{R})} + \|(V_n'-P')w\|_{L^p(\mathbb{R})} \\
&\le CT^{1/4}(a_n)E_{p,n}(w;f') + C\frac{2n}{a_{2n}}T^{1/2}(a_{2n})\|(V_n-P)w\|_{L^p(\mathbb{R})} \\
&\le CT^{1/4}(a_n)E_{p,n}(w;f') \\
&\qquad + C\frac{n}{a_n}T^{1/2}(a_n)(\|(f-V_n)w\|_{L^p(\mathbb{R})} + \|(f-P)w\|_{L^p(\mathbb{R})}) \\
&\le C\left(T^{1/4}(a_n)E_{p,n}(w;f') + T^{3/4}(a_n)E_{p,n}(w;f') + \frac{n}{a_n}T^{1/2}(a_n)\eta\right),
\end{aligned}$$

which shows (4.1). □

[Proof of Theorem 1.1] In (4.1), we take $\eta = C_1T^{1/4}(a_n)E_{p,n}(w;f)$. Then since Lemma 3.5, we have

$$\begin{aligned}
&\|(f'-P')w\|_{L^p(\mathbb{R})} \\
&\le C_2\left(T^{3/4}(a_n)E_{p,n}(w;f') + C_1\frac{n}{a_n}T^{3/4}(a_n)E_{p,n}(w;f)\right) \\
&\le C_2\left(T^{3/4}(a_n)E_{p,n-1}(w;f') + C_1T^{3/4}(a_n)E_{p,n-1}(w;f')\right) \\
&= C_2(1+C_1)T^{3/4}(a_n)E_{p,n-1}(w;f').
\end{aligned}$$

The desired result follows from above. □

Using Theorem 4.1 inductively, we have an estimate in the case of higher order derivative.

Corollary 4.2. Let $w = \exp(-Q)$ be the same as in Proposition 4.1. and $j \in \mathbb{N}$. For any $C_1 \geq 1$, $1 \leq p \leq \infty$ and absolutely continuous function f^{j-1} with $f^(j)w \in L^p(\mathbb{R})$, if

$$\|(f - P)w\|_{L^p(\mathbb{R})} \leq C_1 E_{p,n}(w; f),$$

holds for some $n > j$ and $P \in \mathcal{P}_n$, then there exists constant $C_2 \geq 1$

$$\|w(f^{(i)} - P^{(i)})\|_{L^p(\mathbb{R})} \leq C_2(1 + C_1)T^{(2i+1)/4}(a_n)E_{p,n-i}(w; f^{(i)})$$

holds for all $i = 1, 2, \cdots, j$.

[Proof] We note if $f'w \in L^p(\mathbb{R})$ implies $fw \in L^p(\mathbb{R})$, it follows from $f^{(j)}w \in L^p(\mathbb{R})$ that $f^{(i)}w \in L^p(\mathbb{R})$ for all $0 \leq i \leq j - 1$. We see

$$\|(f' - P')w\|_{L^p(\mathbb{R})} \leq C_2(1 + C_1)T^{3/4}(a_n)E_{p,n-1}(w; f').$$

Applying Theorem 4.1 to $n - 1$, f', P' and

$$\begin{aligned}\eta :&= C_2(1 + C_1)T^{3/4}(a_n)E_{p,n-1}(w; f') \\ &\leq C_2(1 + C_1)T^{3/4}(a_n)\frac{a_{n-1}}{n-1}E_{p,n-2}(w; f''),\end{aligned}$$

we have

$$\begin{aligned}&\|(f'' - P'')w\|_{L^p(\mathbb{R})} \\ &\leq C_2\left(T^{3/4}(a_{n-1})E_{p,n-1}(w; f'') + \frac{n-1}{a_{n-1}}T^{1/2}(a_{n-1})\eta\right) \\ &\leq C_2(1 + C_1)T^{5/4}(a_n)E_{p,n-2}(w; f''),\end{aligned}$$

This shows the case $i = 2$. Repeating this process, we have the result for $1 \leq i \leq j$. $\square$

References

[1] H. N. Mhaskar, Introduction to the Theory of Weighted Polynomial Approximation, World Scientific, Singapore, 1996.

[2] A. L. Levin and D. S. Lubinsky, Orthogonal Polynomials for Exponential Weights, Springer, New York, 2001.

[3] H. S. Jung and R. Sakai, Mean and uniform convergence of Lagrange interpolation with the Erdos-type weights, JIA 2012, 2012:237, doi.10, 1186/1029-242X-2012-237.

[4] R. Sakai and N. Suzuki, On the Favard-type inequalities for exponential weights, Pioneer J. of Math. 3, No. 1 (2011), 1-16.

[5] K. Itoh, R. Sakai and N. Suzuki, The de la Vallée Poussin mean and polynomial approximation for exponential weight, arXiv:1311.3337 [math.CA].

Kentaro Itoh
Department of Mathematics
Meijo University
Tenpaku-ku, Nagoya 468-8502
Aichi Japan
133451501@ccalumni.meijo-u.ac.jp

Ryozi Sakai
Department of Mathematics
Meijo University
Tenpaku-ku, Nagoya 468-8502
Aichi Japan
ryozi@crest.ocn.ne.jp

Noriaki Suzuki
Department of Mathematics
Meijo University
Tenpaku-ku, Nagoya 468-8502
Aichi Japan
suzukin@meijo-u.ac.jp